



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 5, 2017

Mr. Kristian Paul
Regulatory Affairs Manager
Environmental Science Division
Bayer CropScience
2 T.W. Alexander Dr
RTP, NC 27709

Subject: Notification per PRN 98-10 – Minor label changes including moving language, and modifying CA Prop 65 language
Product Name: R & M Garden Dust 5%
EPA Registration Number: 432-1209
Application Date: May 8, 2017
Decision Number: 529728

Dear Mr. Paul:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records. Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 432-1209
Decision No. 529728

If you have any questions, you may contact Eric Bohneblust at (703) 347-0426 or via email at Bohnenblust.eric@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'R. Gebken'.

Richard Gebken
Product Manager 10
Invertebrate & Vertebrate Branch 2
Registration Division (7505P)
Office of Pesticide Programs

R&M Garden Dust 5%

FOR OUTDOOR RESIDENTIAL USE ONLY

- [For Use on Home Garden Vegetables, Fruits, Ornamentals and Lawns]
- [Kills Fleas and Ticks]
- [Kills Japanese Beetles]
- [Kills Over 65 Insects on Vegetables, Fruits, Ornamentals and Lawns]
- [Ready-to-Use Sevin 5% Dust Bug Killer]
- Got Japanese Beetles? Get Sevin!
- Got [insert listed pest here]? Get Sevin!
- Hate The Bugs. Love The Garden. Get Sevin!

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 5.0%

INERT OTHER INGREDIENTS:.....95.0%

TOTAL:100.0%

EPA Reg. No. 432-1209

EPA Est. No. **67572-GA-I**

STOP - KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day X-XXX-XXX-XXXX
For PRODUCT USE Information Call X-XXX-XXX-XXXX

FIRST AID

This product is a N-methyl carbamate and is a cholinesterase inhibitor

If swallowed:	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact X-XXX-XXX-XXXX for emergency treatment information.

NOTE TO PHYSICIAN: CARBARYL is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

NOTIFICATION

432-1209

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

06/05/2017

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

BEE CAUTION: MAY KILL HONEY BEES AND OTHER BEES. This product is highly toxic to honey bees and other bees. Do not apply to plants in bloom.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

~~DIRECTIONS FOR USE~~

Wearing household latex or rubber gloves, use this ready-to-use shaker container to apply product. Dust lightly to cover leaf surfaces with a thin, even film of dust. Apply when air is calm. Start applying at the farthest corner of the treatment area and work backward. For plants above chest height, consider using a liquid carbaryl product.

RESTRICTIONS

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

Non-refillable container. Do not cut package to expose contents. Discard package when empty.

Not for use on plants grown for sale or commercial use, research, or commercial seed production.

PRECAUTIONS

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application. Begin applications when insect or damage is first observed.

HOME GARDEN VEGETABLES AND SMALL FRUITS

Do not repeat applications more than once a week (7 days).

PREHARVEST USE RESTRICTIONS:

Observe the following preharvest intervals (number of days between last application and harvest)

PREHARVEST INTERVAL	
GARDEN CROP	PREHARVEST INTERVAL (DAYS)
Garden beets (roots), radishes, rutabagas, turnips (roots), carrots, potato, blackberries, boysenberries, dewberries, loganberries, raspberries, blueberries, grapes, and strawberries	7
Collards, garden beets (tops), kale lettuce (head and leaf), mustard greens, spinach and turnip (tops)	14
Dried shelled legume beans and peas including: field bean, kidney bean, dry lima bean, navy bean, pinto bean, blackeyed pea, cowpea, Crowder pea, mung bean, southern pea, lentil (dry).	21
Edible-podded legume vegetables including: Snap bean, wax bean, yardlong bean, jack bean, edible-pod pea, snow pea, sugar snap pea, broccoli, brussel sprouts, cabbage, cauliflower, kohlrabi, cantaloupe, cucumber, melons, pumpkins, squash, eggplant, pepper, and tomatoes	3

Vegetables		
CROP	Insect Pests	Comments
Edible-podded legume vegetables including: Snap bean, wax bean, yardlong bean, jack bean, edible-pod pea, snow pea, sugar snap pea. Dried shelled legume vegetables including: field bean, kidney bean, lima bean (dry), navy bean, pinto bean, adzuki bean, blackeyed pea, cowpea, mung bean, Southern pea, lentil (dry). (Do not use on lentils in CA.)*	Bean leaf beetle, cucumber beetles, corn earworm, flea beetles, Japanese beetles, leafhoppers, Lima bean pod borer, Lygus bug, Mexican bean beetle, stink bugs	Maximum number of applications per year: 4
Broccoli, brussel sprouts, cabbage cauliflower, kohlrabi	Armyworms, corn earworm, diamond-back moth larvae, Flea beetles, harlequin bugs imported cabbageworm	Application is only permitted within thirty days from the date of crop emergence or the date of transplanting. Will not control cabbage loopers. Maximum number of applications per year: 4
Collards, garden beets, kale, lettuce, mustard greens, radishes, rutabagas, spinach, turnips	Armyworms, corn earworm, Diamond-back moth larvae, Flea beetles, harlequin bug, Imported cabbageworm, leafhoppers, stink Bugs, tarnished plant bug	Application is only permitted within thirty days from the date of crop emergence or the date of transplanting. Will not control cabbage loopers. Maximum number of applications per year: 4
Cantaloupe, cucumbers, melons, pumpkins, squash	Cucumber beetles, flea beetles, melonworms, pickleworm, squash bug	Maximum number of applications per year: 6
Carrots	Flea beetles, leafhoppers, lygus bugs, spittlebugs, stink bugs	Maximum number of applications per year: 6
Eggplants, pepper, tomato	European corn borer, Fall armyworm, flea beetles, leafhoppers, tomato fruitworm, tomato hornworms, stink bugs,	Maximum number of applications per year: 7
Potato	Colorado potato beetle, flea beetles, leafhoppers, lygus bugs	Maximum number of applications per year: 6

* Not for use on fresh/succulent peas and beans.

SMALL FRUIT		
PLANT	INSECT PESTS	COMMENTS
Blackberries, boysenberries, strawberries, loganberries, dewberries, raspberries	Japanese beetles, leafhoppers, leafrollers raspberry aphid	Maximum number of applications per year: 5
Blueberries	Blueberry maggot, cherry fruitworms, cranberry fruitworms, Japanese beetles	Apply 3 weeks before harvest and repeat 10 days later if necessary. Maximum number of applications per year: 5
Grapes	Grape berry moths, Grape leafroller, grape leafhopper, grape leaf skeletonizer, Japanese beetle, June beetle, red-banded leafroller	Apply just before first brood leafroller larvae emerge from rolls and as needed for leafhoppers. Maximum number of applications per year: 5
Strawberries	meadow spittlebug, strawberry leafroller, strawberry weevil	Maximum number of applications per year: 5

ORNAMENTAL SHRUBS AND FLOWERS

For outdoor use only. To kill bagworm, blister beetles, boxelder bug, boxwood leafminer, flea beetles, Japanese beetle, lacebugs, leafhoppers, leafrollers, periodical cicada, plant bugs, psyllids, rose aphid, rose slug, scale insects in the crawler stage, tent caterpillars and exposed thrips, apply thoroughly to the infested plants at first sign of damage. Do not repeat application more than once weekly or more than six times per year. For best results on scale, apply in spring and early summer when scale crawlers are present.

HOME LAWNS

To kill earwigs, fall armyworm, leafhoppers, millipedes, sod webworm, and sowbugs, sprinkle lightly on grass.

To kill ants (except pharaoh and carpenter ants), chiggers, fleas, spiders (except Brown recluse and Black widow) and ticks, sprinkle lightly on grass. Will only kill pests present at the time of application. Pest must be directly contacted by product to be effective.

For best results, apply after rain or watering and do not water for least 2 days after application. Repeat treatment as needed but not more than every 2 weeks and not more than four times per year.

EARWIGS

This material is effective in killing earwigs and when properly used will reduce the number gaining entrance to homes. Using this ready-to-use shaker container, lightly dust a band 3 to 4 inches wide around the outside foundation wall of the home. Earwigs breed over wide areas around yards and fields and are driven by cold weather to seek shelter in warm buildings. As they move along the foundation wall seeking entrance, they come into contact with the treated barrier and are killed. Be sure to treat in front of all steps and other entrance ways as well as along the foundation wall. Apply also around refuse piles, lumber, mulch and other areas in the yard offering areas of concealment to earwigs. Repeat application in 10 days or when earwigs become troublesome. If the dust barrier is washed away by rain, it should be replaced with a new application if earwigs are still a problem.

This container is sold by weight, NOT by volume. You can be assured of the proper weight even though some settling of contents normally occurs during transportation and handling.

Important: Read the entire Directions before using this product. Disclaimer: Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions for Use. To the extent consistent with applicable law, manufacturer makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty. To the extent consistent with applicable law, in no case shall Manufacturer be liable for consequential, special or indirect damages resulting from the use, handling or application of this product.

To obtain additional information or in case of an emergency call 770-787-1977.

- If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call: (800) 420-9347.
- If you have questions relating to product use or application, please call (800) 969-7200.
- If you have a product spill or transportation emergency call (800) 424-9300.
- For Material Safety Data Sheets please call (800) 969-7200.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store pesticide in original container and place in a locked storage area.

Pesticide Disposal:

If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-XXX-XXX-XXXX for disposal instructions. Never place unused product down any indoor or outdoor drain.

[Optional language (only if required by the State of CA):

~~**WARNING:** This product contains carbaryl, a chemical known to the State of California to cause [cancer] and [reproductive harm] in laboratory animals.]~~  **NOTICE:** Cancer – www.P65Warnings.ca.gov

SEVIN is a registered trademark of Tessenderlo Kerley, Inc.

PRODUCED FOR



Bayer Environmental Science

A **Division Business Group** of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

R&M Garden Dust 5% (Pending) 05/05/2017